SMALL ODONTOCETES

(SHEET 2)



Southern right whale dolphin
(Lissodelphis peronii)

- specimen
- specimen, exact locality not known
- specimen now thought to have been incorrectly identified as *L. peronii* (P.J.H. van Bree, personal communication)
- stranding observation
- at-sea observation
- observation, exact locality not known
- observations within this area
- numerous observations along ship track
- scattered observations along ship track

NORTHERN RECORDS

Known from two observations north of the region mapped: 29°35′S, 71°45′W and 18°53′S, 71°45′W (ref. 9).

Note

Localities from references 6, 23, 31, 41, 50, 54, 60, 72, 79, and 88 are plotted as shown in ref. 31.



Long-finned pilot whale
(Globicephala melaena)

- specimen
- at-sea observation
- observation, exact locality not known
- seen frequently



Beaked whales

- beaked whale, species not known

Southern bottlenose whale
(Hyperoodon planifrons)

- specimen
- specimen, exact locality not known
- catch record
- at-sea observation
- at-sea observation, identification uncertain

Arnoux's beaked whale
(Berardius arnuxii)

- specimen
- stranding observation
- at-sea observation

NORTHERN RECORDS

H. planifrons, one observation and two strandings: observation off the coast of Chile at 33°S, 73°27′W (ref. 17); stranding (holotype specimen) at Lewis Island, northwestern Australia (ref. 27); stranding on the coast of the province of Buenos Aires (ref. 61). *B. arnuxii*, one stranding: south of La Plata, Argentina (ref. 71).

Killer whale
(Orcinus orca)

- specimen
- stranding observation
- at-sea observation
- seen frequently
- scattered observations along ship track

NOTES

(1) Numbers by symbols refer to bibliography
(2) Maps compiled by Robert L. Brownell, Jr., drawings by Alice R. Tangerini and Bernardo Menendez

AMERICAN GEOGRAPHICAL SOCIETY—1974