

MAP 1

- **Anasterias**
  - 1 *antarctica* (0–120 m)
  - 2 *conferta* (0–7 m)
  - 3 *directa* (intertidal)
  - 4 *mawsoni* (littoral)
  - 5 *minuta* (0–16 m)
  - 6 *pedicellaris* (7–95 m)
  - 7 *perrieri* (0–22 m)
  - 8 *rupicola* (intertidal)
  - 9 *sphaerulata* (intertidal)
  - 10 *stolidota*
  - 11 *studeri* (50–278 m)
- **Anteliaster**
  - 12 *australis* (79–341 m)
  - 13 *scaber* (22–113 m)
- **Asterodon**
  - 14 *dilatatus* (4–100 m)
  - 15 *miliaris* (35–100 m)
  - 16 *robustus* (intertidal)
  - 17 *singularis* (shore–78 m)
- **Crossaster**
  - 18 *australis* (120 m)
  - 19 *canopus* (322–340 m)
  - 20 *japonicus* (90–600 m)
  - 21 *penicillatus* (55–280 m)
- **Notasterias**
  - 22 *armata* (15–752 m)
  - 23 *bongraini* (60–830 m)
  - 24 *haswelli* (60–354 m)
  - 25 *pedicellaris* (3100 m)
  - 26 *stolophora* (247–274 m)
- **Paralophaster**
  - 27 *asperatus* (110 m)
  - 28 *godfroyi* (193–567 m)
  - 29 *lorioli* (1375–5500 m)
- **Pergamaster**
  - 30 *incertus* (247–540 m)
  - 31 *synaptorus* (342 m)
  - 32 *tessellatus* (110–1700 m)
  - 33 *triseriatus* (369–384 m)
- **Saliasterias**
  - 34 *brachiata* (15–540 m)



MAP 2

- **Coscinasterias**
  - 1 *africana* (0–40 m)
  - 2 *calamaria* (4–80 m)
  - 3 *candicans* (450–560 m)
  - 4 *eustyla* (200–300 m)
  - 5 *victoriae* (70 m)
- **Cosmasterias**
  - 6 *capensis* (120 m)
  - 7 *dyscrita* (230–400 m)
  - 8 *lurida* (0–690 m)
  - 9 *polygramma* (490 m)
  - 10 *tomidata* (90 m)
- **Diplopteraster**
  - 11 *perigrinator* (254 m)
  - 12 *semireticulatus* (99–138 m)
  - 13 *verrucosus* (75–205 m)
- **Patiriella**
  - 14 *calcar* (intertidal)
  - 15 *exigua* (intertidal)
  - 16 *fimbriata* (2–360 m)
  - 17 *gunnii* (intertidal)
  - 18 *regularis* (0–100 m)
- **Gaussaster**
  - 19 *antarcticus* (300 m)
  - 20 *vanhoeffeni* (2450–2725 m)
- **Kampylaster**
  - 21 *incurvatus* (93–472 m)
- **Luidiaster**
  - 22 *antarcticus* (3900 m)
  - 23 *gerlachei* (160–810 m)
- **Rhopiella**
  - 24 *hirsuta* (151–603 m)
  - 25 *koehleri* (93–130 m)
  - 26 *pterasteroides* (358 m)
  - 27 *smilax* (110–350 m)
  - 28 *smithi* (450 m)



Coscinasterias
Arm-spread up to 14 in.



Patiriella
Arm-spread up to 3 in.

Cosmasterias
Arm-spread up to 8 in.

All drawings after Fell, 1962



Asterodon
Arm-spread up to 5 in.

Crossaster
Arm-spread up to 7 in.



MAP 3

- **Cuenotaster**
  - 1 *involutus* (60–578 m)
- **Granaster**
  - 2 *nutrix* (1–250 m)
- **Lysasterias**
  - 3 *adeliae* (24–578 m)
  - 4 *belgicae* (560 m)
  - 5 *chirophora* (450–560 m)
  - 6 *digitata* (163–300 m)
  - 7 *hemiora* (70 m)
  - 8 *heteractis* (342 m)
  - 9 *joffrei* (88–810 m)
  - 10 *perrieri* (0–708 m)
- **Myoraster**
  - 11 *antarcticus* (88–578 m)
  - 12 *incertus* (littoral)
  - 13 *tenuis* (110 m)
- **Pedicellaster**
  - 14 *formatus* (120 m)
  - 15 *hypernotius* (93–450 m)
  - 16 *reticulatus* (156 m)
  - 17 *scaber* (40–50 m)
- **Perknaster**
  - 18 *aurantiacus* (75–254 m)
  - 19 *aurorae* (18–300 m)
  - 20 *charcoti* (132–235 m)
  - 21 *densus* (101–385 m)
  - 22 *fuscus* (3–362 m)
  - 23 *rallieri* (littoral)
  - 24 *sladeni* (120–500 m)



MAP 4

- **Anareaster**
  - 1 *gourdoni* (10–540 m)
- **Cryptasterias**
  - 2 *brachiata* (25–30 m)
  - 3 *turqueti* (0–45 m)
- **Neosmilaster**
  - 4 *georgianus* (1–335 m)
  - 5 *steineni* (100–160 m)
- **Odinella**
  - 6 *nutrix* (125–640 m)
- **Psalidaster**
  - 7 *mordax* (163–367 m)

COMPILED BY H. B. FELL
AND S. DAWSEY

AMERICAN GEOGRAPHICAL SOCIETY—1969